



Required Immunizations and Tests for Students

The Corvallis Clinic requires the following tests and vaccines for any health care personnel (HCP). HCP refers to all paid and **unpaid** persons working in the healthcare setting who have the potential for exposure to patients and/or infectious materials. This includes students who are completing a rotation in the healthcare setting.

Hepatitis B

Provide one of the following as evidence of Hepatitis B (HBV) immunity:

- Three Hepatitis B vaccines OR
- Laboratory evidence of Hepatitis B immunity (titer*)

TB Screening

Confirmation of a negative tuberculosis (TB) test is required within 12 months.

If you have no record of a TB test, you will need to have one taken. This can be done at TCC Occupational Medicine Department, or the facility of your choice.

If you've had a positive TB test in the past, you will need to complete a TB Risk Factor Screening form and chest x-ray or provide TCC with a record of a chest x-ray dated within the last 6 months.

Measles (Rubeola), Mumps & Rubella

Provide one of the following as evidence of MMR immunity:

- Two MMR vaccines OR
- Two Measles (Rubeola), two Mumps and one Rubella vaccine OR
- Laboratory evidence of Measles, Mumps and Rubella immunity (titer*)

Varicella (Chicken Pox)

Provide one of the following as evidence of Varicella immunity:

- Two varicella vaccines OR
- If you had chicken pox as a child, provide laboratory evidence of Varicella immunity (titer*)

Influenza

Influenza is a serious disease that can lead to hospitalization and sometimes even death. Annual influenza vaccination is highly encouraged for all healthcare workers and students.

Tdap

Provide evidence of a single dose of the Tetanus, Diphtheria, and Pertussis (Tdap) vaccine.

Reviewed 4/7/22

Adapted as per TCC Policy #230-00-022, Employee Vaccination and Health Screening & updated to align with Oregon Rule 333-019-1010



Required Immunizations and Tests for Students

COVID-19 Vaccination

Provide proof of full vaccination status. 2 weeks following second dose in a 2-dose series, or 2 weeks following a single dose vaccine.

***Definition of a titer:** A titer is a laboratory test that measures the level of antibodies in a blood sample to determine immunity.

Reviewed 4/7/22

Adapted as per TCC Policy #230-00-022, Employee Vaccination and Health Screening & updated to align with Oregon Rule 333-019-1010